

Beta cell connectivity in pancreatic islets

Rutter, Guy A; Hodson, David J

DOI:

[10.1007/s00018-014-1755-4](https://doi.org/10.1007/s00018-014-1755-4)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Rutter, GA & Hodson, DJ 2015, 'Beta cell connectivity in pancreatic islets: a type 2 diabetes target?', *Cellular and Molecular Life Sciences*, vol. 72, no. 3, pp. 453-67. <https://doi.org/10.1007/s00018-014-1755-4>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

The final publication is available at Springer via <http://dx.doi.org/10.1007/s00018-014-1755-4>

Checked December 2015

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

REVIEW

Beta cell connectivity in pancreatic islets: a type 2 diabetes target?

Guy A. Rutter and David J. Hodson

Section of Cell Biology, Department of Medicine, Imperial College London, Imperial Centre for Translational and Experimental Medicine, Hammersmith Hospital, Du Cane Road, London W12 0NN, UK.

Correspondence

Dr David J. Hodson
Section Cell Biology
Department of Medicine
Imperial College London
London SW7 2AZ
United Kingdom
e-mail: d.hodson@imperial.ac.uk
Tel: +44 (0)20 7594 1713
Fax: +44 (0)20 7594 3351

Prof. Guy A. Rutter
Section of Cell Biology
Department of Medicine
Imperial College London
London SW7 2AZ
United Kingdom
e-mail: g.rutter@imperial.ac.uk
Tel: +44 (0)20 7594 3340
Fax: +44 (0)20 7594 3351

Summary sentence: beta cell communication during type 2 diabetes

Pubmed was searched using combinations of the following: “islet”, “diabetes”, “electrical”, “activity”, “channels”, “calcium”, “sodium”, “potassium”, “incretin”, “GLP-1”, “GIP”, “coordination”, “synchrony”, “dynamics”, “gap junction”, “connexin 36”, “paracrine”, “architecture”, “structure”, “autocrine”, “genetics”, “neural”, “tcf7l2”, “adcy5” and “gipr”.

DISCLOSURE STATEMENT: The authors have nothing to disclose.

Word count: 4746 (main text)

Keywords: mouse, human, signaling, insulin, diabetes, imaging, network

Abstract

Beta cell connectivity describes the phenomenon whereby the islet context improves insulin secretion by providing a three-dimensional platform for intercellular signaling processes. Thus, the precise flow of information through homotypically interconnected beta cells leads to the large-scale organization of hormone-release activities, influencing cell responses to glucose and other secretagogues. Although a phenomenon whose importance has arguably been underappreciated in islet biology until recently, a growing number of studies suggest that such cell-cell communication is a fundamental property of this micro-organ. Hence, connectivity may plausibly be targeted by both environmental and genetic factors in type 2 diabetes mellitus (T2DM) to perturb normal beta cell function and insulin release. Here, we review the mechanisms that contribute to beta cell connectivity, discuss how these may fail during T2DM, and examine approaches to restore insulin secretion by boosting cell communication.

Abbreviations: AC, adenylyl cyclase; ACh, acetyl-choline; ADP, adenosine diphosphate; ATP, adenosine triphosphate; cAMP, cyclic adenosine monophosphate; Cx36, connexin 36; Epac, exchange protein activated by cAMP; fMCI, functional multicellular calcium imaging; GABA, gamma aminobutyric acid; GIP, glucose-dependent insulintropic polypeptide; GJ, gap junction; GLP-1, glucagon-like peptide-1; GWAS, genome-wide association studies; GPCR, G-protein coupled receptor; K_{ATP} , ATP-sensitive K^+ channel; SST, somatostatin; SNP, single nucleotide polymorphism; T2DM, Type 2 diabetes mellitus; VDCC, voltage-dependent Ca^{2+} -channel.

1 Introduction

2 Type 2 diabetes mellitus (T2DM) is a global epidemic that currently consumes ~10% of the
3 healthcare budget in the developed world [1]. This syndrome has a complex aetiology but can be
4 summarised as a failure of the beta cell mass to adequately compensate for insulin resistance, or
5 alternatively a primary beta cell defect that leads to insulin resistance. The resulting glucose
6 intolerance, coupled with dyslipidemia, drives a range of costly secondary complications
7 including retinopathy, vasculopathy, renal failure, cancer and cardiovascular disease [2,3].
8 Consequently, elucidation of the mechanisms underlying the control of insulin secretion from
9 individual beta cells has been the focus of intense research efforts. Thus, in response to an
10 elevation of blood glucose, equilibration of the sugar across the plasma membrane occurs rapidly
11 and is achieved *via* either the low affinity glucose transporter Glut2/*slc2a2* (rodents) or the higher
12 affinity transporter Glut1/*slc2a1* (man) [4]. The low affinity hexokinase, glucokinase is then
13 chiefly responsible for determining glycolytic flux towards pyruvate [5]. Conversion of the latter
14 to acetyl-CoA in the mitochondrial matrix, and its oxidation via the tricarboxylate cycle, then
15 ensues [6,7]. The resultant increases in the ratio of free ATP to ADP (ATP:ADP) in the cytosol
16 [8] and sub-plasma membrane domain [9] then leads to closure of ATP-sensitive K⁺ channels
17 (K_{ATP}), membrane depolarisation and the influx of calcium (Ca²⁺) through voltage-dependent
18 Ca²⁺-channels (VDCC) [6,7,10,11]. Together with the activation of a less well-defined
19 “amplifying” pathway [12,13], localized increases in the intracellular free Ca²⁺ concentration
20 [14], including at the surface of the secretory granule [15], then provoke insulin release through
21 interactions with the exocytotic machinery [16,17].

22
23 By comparison, the population-level regulation of insulin release is less well understood,
24 although the idea that it may contribute to T2DM risk has been suggested [18-22]. Providing
25 evidence that cell-cell interactions are a prerequisite for proper hormone secretion is the
26 observation that beta cells *incommunicado* (*i.e.* as isolated cells) release less insulin *per capita*
27 than their properly-connected counterparts within the intact islet [19,23,24]. Indeed, a feature of
28 the endocrine pancreas is the three-dimensional encapsulation of beta, and other cell types, into
29 islets of Langerhans, a biological scaffold for cell-cell communications. Since these microorgans
30 are conserved throughout the mammalian kingdom and beyond [25], albeit with important
31 differences in the numbers of each cell type and their arrangement within the islet (see below),
32 the intraislet mechanisms governing insulin secretion may represent an underappreciated target
33 through which T2DM insults provoke hyperglycemia. Building upon recent findings from our
34 own [26-28] and others’ [20-22,29-31] laboratories, the aim of the present review is to describe
35 our current understanding as to how beta cell-beta cell communication (hereafter referred to as

“connectivity”) contributes to the normal regulation of insulin secretion in healthy subjects. We also discuss how changes in this property may contribute to T2DM risk in genetically-susceptible individuals.

Islets as discrete secretory units

The term “endocrine pancreas” describes the thousands (millions in man) of islets of Langerhans scattered throughout the exocrine tissue. Each islet can range in size from 20-400 μm and comprises alpha- (glucagon), beta- (insulin), delta- (somatostatin), epsilon- (ghrelin) and pancreatic polypeptide (PP) cells. Strikingly, islets are evolutionarily-stable structures and are present in most mammals studied to date, including the Beluga whale, with a similar range of sizes reported in each species [25]. With the exception of bats, horses, hyenas, primates and humans, the arrangement of endocrine cells within islets is similar [25]. Thus, in rodent islets, the most-studied model, beta cells form a central core, with alpha cells occupying the mantle [25,32,33]. Suggesting that this may be a consequence of the vasculature, blood flow has been shown to follow an inner-outer flow pattern, irrigating beta before alpha cells in this species [34], and the vasculature appears to be instructive for pancreas development [35]. By contrast, beta cells in human islets are interspersed with alpha cells, in part the consequence of the tertiary folding of an initial trilaminar alpha-beta-alpha sheet, which promotes heterologous contacts [33,36-38]. As well as differences in islet architecture, alterations to cell proportion are also apparent between species. For example, the ratio of beta:alpha cells in rodent islets is ~4:1, whereas in humans it is ~1.25:1. Such divergence in islet architecture likely influences cell-cell communication by altering the extent and nature of cell-cell signaling processes, and may be an important source of species differences in islet function. Regardless, the islet structure is permissive for insulin secretion, and beta cells in two dimensions display blunted responses to input, both in terms of Ca^{2+} signaling and magnitude hormone release [27,39-41].

High-speed imaging of beta cell connectivity

Over the last decade, advances in microscopy have allowed cell dynamics to be monitored *in situ* within the intact tissue setting [42]. Key to this is the use of high-speed imaging, which when combined with highly-sensitive detectors, allows a large area to be rapidly traversed at cellular resolution. In terms of endocrine organ function, the physiologically relevant output is hormone release. However, large-scale imaging of exocytosis in individual cells is only just becoming possible, although the currently available dyes possess signal-to-noise ratios incompatible with high-speed acquisition at visible light wavelengths [43-46]. To circumvent these issues, membrane voltage or intracellular Ca^{2+} concentrations can instead be used as a proxy for Ca^{2+} -

dependent hormone release [47-50]. To this end, functional multicellular Ca^{2+} imaging (fMCI), originally used to map activity in cortical circuits [51-53], has recently been adapted for use in beta cells [27,28]. By coupling a laser bank to a Nipkow spinning disk, the millisecond organization of beta cell population Ca^{2+} -spiking activity can be captured in near real-time with reduced phototoxicity and photobleaching. Following acquisition, the datasets are subjected to non-deterministic Monte Carlo-based models to identify the cells with similar behavioural profiles, *i.e.* those with correlated activity and which are assumed to contribute to the same secretory process [42,54]. Statistical significance is determined by shuffling the experimental dataset and calculating the likelihood of detecting the same correlation pattern due to chance. A functional connectivity map can then be constructed based on the location of significantly correlated cells pairs, allowing perturbations to beta cell connectivity to be evaluated (see Figure 1, top panel, for an example). In a refinement of this method, beta cell metabolic interconnectivity has recently been mapped in intact islets by monitoring intracellular free ATP:ADP dynamics, as for Ca^{2+} [55]. When using these techniques, it is important to note that the territories of communicating beta cells within intact islets are larger than those that can be recorded, limiting any physiological inferences that can be drawn.

Islet wiring patterns

Network science principally relies on the use of graph theory to identify the interactions that govern behavior in complex systems (see [42] for a review of network science in Endocrinology). Using these approaches, it has become increasingly clear that network topology tends to be conserved (*e.g.* scale-free and random) irrespective of the components examined (*e.g.* cells *versus* people) [42,56,57]. Recent research has shown that graph theory is also applicable to the description of complex dynamics in the endocrine pancreas. Thus, analysis reveals that beta cells comprise glucose-responsive scale-free networks in which cells can communicate over long distances, through presently undefined mechanisms [29]. Such network topologies are defined by a power-law distributed link probability in which a minority of cells (termed highly-connected nodes) host the majority of connections and are said to possess small-world properties if there is a tendency towards formation of cliques (six degrees of separation concept) (Figure 1, bottom panel). Price was the first to describe scale-free networks, noting that journal citations follow a power-law distribution, sharing features in-keeping with Pareto's law (the 'rich-get-richer' hypothesis) [58]. Subsequently, Barabasi and Albert showed that preferential attachment is responsible for the emergence of scale-free properties [59]. Notably, scale-free distributions are ubiquitous and have been described in social networks, computer networks, neural networks and anterior pituitary networks [54,60-63]. An important feature of scale-free networks is robustness

at low wiring cost: the chances of a random attack disabling communication are low and the use of hubs to route information reduces signal transmission length [42]. However, should the highly-connected nodes be specifically targeted, the network is vulnerable to collapse, since a high proportion of links will be lost (Figure 1, bottom panel). Therefore, an interesting but untested possibility is that highly-connected beta cell nodes may represent a subpopulation which is particularly susceptible to T2DM insults. Conversely, these highly-connected nodes may serve as a functional reserve to maintain islet function in the face of gross perturbation by allowing the redistribution of information, again, a hypothesis that requires experimental validation.

Mechanisms underlying beta cell-beta cell connectivity

Neural circuits have a clear basis for long-range connectivity, since neurons send out axonal projections that can form synapses located millimetres apart. By contrast, it is less easy to conceptualize how beta cells within the islet can communicate over long distances to organize their activities. Might this involve, for example, “physical connections” (*e.g.* through islet interneurons) between remote cells, or alternatively linearly-connected “trains” of beta or other cells along which signals are transmitted to a distant cell(s) from a controller (“pacemaker”) at a coordination hub? In any case, the islet possesses a formidable signaling toolbox (see Figure 2). This is reviewed in depth elsewhere [20,21,28,64], so here we limit our discussion to the pathways which may conceivably underlie connectivity between beta cells.

Gap junctions: The best characterized cell-cell coupling mechanism in the pancreas is provided by gap junctions (GJ). Beta cells within rodent and human islets are homotypically-connected by connexin 36 (Cx36 or GJD2) [65,66]. GJs comprised of Cx36 are charge and size-selective channels that allow the intercellular passage of ions (*e.g.* Ca^{2+} , Na^{+} and Zn^{2+}) and nucleotides (*e.g.* ATP) [19,20,67]. Providing evidence that Cx36 is critical for coordinating islet activity are the observations that dispersed beta cells fail to synchronize their responses to glucose, and islets lacking Cx36 display more stochastic activity patterns due to increases in beta cell functional heterogeneity [31,68-70]. GJ-linkages are essential for the regulation of normal hormone release, since mice deleted for Cx36 are glucose intolerant and display impaired pulsatility, as well as elevated basal insulin secretion [22,68,71]. It is unclear how GJs could account for the long-range functional connections that project between distant cells, as practically all beta cells express Cx36 protein, meaning that communication should encompass even close neighbors [65,72]. However, heterogeneity exists in fluorescence recovery after photobleaching (FRAP) within islets [73], suggesting that connectivity patterns between individual beta cells may at least reflect differences in functional GJ coupling. As proposed above, this may lead to the formation of linear groups of

cells, tightly interconnected in three dimensions between one another, but (relatively) isolated from neighboring cells outside the train, thus forming a conduit for the passage of ionic (Ca^{2+}) or other (*e.g.* paracrine, see below) signals.

Neural: Islets receive rich innervation from the autonomic nervous system, and neural regulation of insulin secretion is critical for normal glucose homeostasis *in vivo*. The existence of a physical network of neurons to couple remote beta cells within the islet thus provides a conceptually straightforward model to explain recent experimental observations [26,27,29]. Indeed, insulin release is strongly stimulated by postganglionic cholinergic fibres that signal via acetylcholine (ACh)-mediated activation of muscarinic receptors to phase-set and synchronize beta cell activity within and, potentially, between islets [74-76]. Such activation underpins the cephalic phase of insulin secretion in anticipation of food [77]. In addition, other neuropeptides including pituitary adenylate cyclase activating peptide (PACAP) and vasoactive intestinal peptide (VIP) may contribute to the parasympathetic control of beta cell function [74,78]. By contrast, insulin release is suppressed by noradrenergic sympathetic neurons that signal via α 2-adrenoreceptors to open K_{ATP} channels [74,79,80], although a stimulatory effect of noradrenaline has also been observed, probably through effects on cAMP accumulation and β -adrenoreceptor activation [81,82]. Marked differences exist in the neural regulation of insulin secretion between rodents and man. Thus, human islets are relatively devoid of parasympathetic nerve fibres [83], and glucose-sensitization of beta cell activity instead relies upon ACh release from vesicular acetylcholine transporter-expressing alpha cells [84,85]. This lack of direct innervation may partly explain why beta cell glucose responses in human islets are largely stochastic, with synchrony detected only between small cell clusters [27,33,86]. Conversely, the assessment of whether neurons contribute to long-range connectivity in mouse islets firstly requires confirmation of cholinergic fibre survival in isolated islets, followed by their specific manipulation (*e.g.* using patch clamp).

Primary cilia: Cilia can be regarded as cell extensions that act as signaling hubs due to expression of G-protein coupled receptors, ion channels and transcription factors [87]. Primary cilia are immotile and are formed from a ring of nine microtubule doublets wrapped in a membrane sheath [88]. While studies of *Kif3a*, *Lkb1* and *Rfx3* knockout mice have all invoked a role for cilia in pancreatic development (*i.e.* ductal and endocrine cell specification) [87,89-91], little is known about their involvement in cell-cell signaling processes within the islet. Given the role of cilia in signal transmission in other tissues [92], and potentially in exosome-mediated intercellular communications [93], we believe this warrants further investigation.

Paracrine signaling:

Intercellular communication may also be possible *via* the production and secretion of messengers which act on neighboring cells [20,21,28]. Over 230 secreted factors have been identified in rodent islets [94], and a number of signalling loops with roles in the regulation of beta cell function and insulin release are now well characterised (see references [21,28,64]). Despite this, it is unclear how paracrine factors could contribute to the complex functional islet wiring patterns described using graph theory [29,30], since all beta cells within the molecule diffusion path would be expected to be affected. Although it is plausible that active transport mechanisms and cognate receptor expression levels/patterns may allow more precise communication between beta cells, this needs further study.

Despite the plethora of signaling mechanisms available within the islet, we suggest that a combination of modalities is required for producing the complex activity patterns that underlie beta cell-beta cell communication and connectivity. Notably, differences in signaling input, together with alterations to islet architecture, may play an important role in determining species-specific responses to secretagogues such as glucose and incretins.

Glucose and GLP-1-regulated connectivity: metabolic signals

It is generally acknowledged that metabolic activity within individual beta cells is oscillatory, and that this generates the membrane bursting activity required for Ca^{2+} influx and exocytosis [95]. Whether metabolic oscillations are driven by Ca^{2+} oscillations, or *vice versa*, is still the source of debate [95,96], but the islet context seems to be critical, since dispersed beta cells display reduced periodicity in mitochondrial potential [97]. Moreover, total internal reflection fluorescence (TIRF) microscopy of mouse islets has shown that near-membrane glucose-induced oscillations in ATP:ADP are coordinated between small beta cell clusters [98], confirming earlier observations that employed lower resolution autofluorescence imaging of NAD(P)H [99-101]. The mechanisms underlying the synchronous propagation of energy status between beta cells remain unknown, but may reflect Ca^{2+} feedback and intrinsic metabolic behaviour [96], or alternatively, metabolic coupling *via* GJs [102,103].

In addition to glucose, secretory potentiators, including members of the incretin family, are able to influence beta cell energetics. The incretin, glucagon-like peptide 1 (GLP-1), is released from the gut in response to bile transit and glucose-dependently augments insulin secretion [104-106]. While its effects on cAMP-Epac2, MAPK and beta-arrestin signaling pathways are well-characterised [107-109], little is known about whether GLP-1 alters the beta cell metabolic

setpoint to influence ATP:ADP. Whereas luciferase-based studies by us have demonstrated a role for GLP-1 in mitochondrial ATP synthesis in clonal MIN6 beta cells [110], others have observed no effect of the incretin in rodent islets using biochemical detection methods [111]. Since ATP dynamics and/or cell heterogeneity may mask actions of incretin on metabolism, the effects of GLP-1 on intracellular free ATP:ADP were monitored with cellular resolution by expressing the recombinant probe Perceval throughout the first few layers of rodent and human islets [8,55,112]. Using these methods, we found that GLP-1 engages a metabolically-coupled subnetwork of beta cells to amplify insulin secretion, an action that is dependent upon Ca^{2+} influx and elevations in cAMP [55]. Of note, in these studies, beta cells within mouse islets responded coordinately to GLP-1 with synchronous ATP:ADP oscillations, whereas human islets exhibited more random dynamics. Thus, the regulation of beta cell-beta cell metabolic connectivity may potentially contribute to the disparate actions of incretin in rodents and man, although confirmation of this will require simultaneous measures of Ca^{2+} and ATP:ADP in islets of both species.

Glucose- and GLP-1-regulated connectivity: Ca^{2+} signals

Ca^{2+} -imaging of pancreatic islet slices has revealed that glucose likely drives large-scale increases in population synchrony by coaxing activity in a scale-free and small-world network of beta cells [29,30,49]. Notably, propagation of Ca^{2+} waves *via* GJs is hypothesised to underlie islet dynamics in response to glucose, since the length of individual correlated links depends on Euclidean distance, although long-range communications are still evident [29]. Confirming these findings, we have recently shown that the rapid (ms) oscillations in electrical activity are similarly dictated by scale-free and small-world beta cell wiring patterns [113]. Thus, under conditions of high glucose, beta cells work together as defined subpopulations to orchestrate and drive insulin release from the islet.

As well as glucose, insulin secretion is also reliant upon the amplifying or potentiating actions of incretins. Indeed, in humans, almost 70% of the insulin-raising effects of oral glucose challenge can be attributed to the incretin effect [114]. Notably, the insulinotropic activity of exogenously-administered GIP and GLP-1 is diminished in T2DM [115,116], suggesting that altered beta cell incretin responsiveness may contribute to the disease state, although causality is not well defined [117]. Since the single biggest T2DM risk factor remains obesity, and high BMI individuals present with reduced GLP-1-stimulated insulin secretion [118,119], excess lipid may target incretin action to impair beta cell function. To investigate this, we subjected human islets to fMCI to map population dynamics, and found that both GIP and GLP-1 recruit a highly coordinated subnetwork of GJ-coupled beta cells to augment insulin secretion [27,28]. This process of

1 incretin-regulated beta cell connectivity may be a target for the insulin-lowering effects of free
2 fatty acid (FFA), since it could be disrupted in a GJ-dependent manner following exposure to a
3 lipotoxic milieu, and was inversely correlated with donor BMI [27]. Mechanistically, this may
4 involve FFA-induced overexpression of inducible cAMP early repressor gamma (ICER- γ), a
5 protein that binds a cAMP-response element in the Cx36 promoter [120,121]. By contrast, a
6 similar effect of incretin on beta cell interactivity was not present in mouse islets, but could be
7 revealed by placing mice on a high fat diet to disrupt normal glucose responses [27,28]. We
8 therefore speculate that such divergent regulation of the incretin axis, potentially stemming from
9 structural and functional differences in islet architecture, may represent a novel target for pro-
10 diabetogenic insults in man.

11 12 **Genes and connectivity**

13 Type 2 diabetes has a strong hereditary component [122-124]. Consequently, genome wide
14 association studies (GWAS) have identified a number of gene variants linked with an increased
15 odds ratio (OR) of developing elevated fasting glucose and T2DM. Although the effects of these
16 variants are usually quite small, their very existence indicates that genes in the associated loci are
17 highly likely to play a role in disease aetiology [125,126]. While gene variants and glucose
18 homeostasis are well studied in man, relatively less is known about their precise mechanisms of
19 action at the islet level [125], and in particular upon beta cell connectivity. Several dozen (>90)
20 risk-associated polymorphisms have been identified to date, and those with the strongest OR for
21 development of T2DM, or with known effects on beta cell-cell communication, are discussed
22 below (see Figure 3):

23
24 ***TCF7L2***: *TCF7L2* is a member of the canonical Wnt-signaling pathway and a transcriptional
25 partner for beta-catenin. Individuals who possess a single nucleotide polymorphism (SNP),
26 rs7903146, in intron 3 of the *TCFL72* gene on chromosome 10, have an increased risk of
27 developing T2DM, with an OR of 1.45 for the T allele [127-130]. This is believed largely to be
28 due to defects in insulin secretion (insulin sensitivity is slightly impaired in T allele carriers), as
29 well as a markedly (~50%) attenuated incretin effect [127,131,132] (though see [125] for a
30 discussion of a role for hepatic glucose handling). Although the subject of debate, these results
31 have subsequently been confirmed in conditional rodent models and human islets. Thus, *TCF7L2*
32 silencing leads to impaired insulin secretion from isolated mouse and human islets [133,134], and
33 deletion of *Tcf7l2* throughout the pancreas or selectively in the beta cell causes glucose
34 intolerance [135,136], particularly after oral glucose administration, with the observed effects
35 increasing with age or exposure to a high fat diet (HFD). Of note, a further study failed to detect

any effects on glycemia of deleting *Tcf7l2* in the adult beta cell, although this report was restricted to examination of intraperitoneal glucose tolerance in young (<12 wks) animals [137]. GLP-1-stimulated insulin secretion is strongly inhibited by *Tcf7l2* elimination *in vitro* [134,135], the latter due largely to reduced GLP-1R expression and defects in the exocytotic apparatus [133,135,138,139]. Interestingly, when investigated in dissociated islets, TCF7L2 knockdown leads to a slight potentiation of glucose-induced Ca^{2+} increases [133,140], although only single (or clusters) of beta cells were studied, precluding analysis of synchrony or coordination. By contrast, ablation of the *Tcf7l2* gene selectively in the beta cell through *Ins1Cre*-directed recombination of *lox*'d alleles impairs these increases when assessed in the intact islet setting [136]. The reasons for these differences remain obscure but suggest that either silencing in non-beta cells in the former case, or altered beta cell-beta cell interactions in the latter, are at play. Of note, *Tcf7l2* silencing in INS1 cells lowers the expression of Ca^{2+} channel subunits [141], suggesting that TCF7L2 may exert control, either directly or indirectly, over the Ca^{2+} -signaling machinery. Of relevance, when studied in islets from mice maintained on a high fat diet (HFD), glucose-stimulated beta cell connectivity in *Tcf7l2* null animals was significantly reduced *versus* that of control animals [136] (manuscript submitted). Of note, this alteration was not associated with any changes in GJ mRNA expression, though may conceivably involve changes in Cx36 protein abundance.

ADCY5: ADCY5 gene products encode isoform V of the adenylate cyclase family, a type III Ca^{2+} -inhibited enzyme tasked with generation of cAMP [142,143], a second messenger involved in glucoregulation as part of the “amplifying” pathway [144]. Whereas other isoforms predominate in the rodent islet, ADCY5 is amongst the most abundant members of this family in human beta cells [26,145]. The T2DM-associated SNP rs11708067 on chromosome 3 lies within intron 3 of the *ADCY5* gene and is associated with increased fasting glucose and 2-hour glucose, but not oral glucose responses [146], with an OR of 1.23 for the major A-allele [147]. Using lentiviral shRNA approaches to silence gene and protein expression in human islets, we have recently shown that ADCY5 is required for the coupling of glucose but not incretin to insulin secretion [148]. Although the former is partly due to impaired insulin processing (*i.e.* proinsulin → insulin conversion) [149], islets depleted for ADCY5 also displayed impaired glucose- but not GLP-1-induced increases in cAMP, and consequent impairments in glucose-induced metabolism (ATP:ADP ratios). Moreover, ADCY5-silenced islets showed more stochastic long-term evolutions in coordinated beta cell activity following glucose exposure [148]. By contrast, GLP-1-regulated connectivity was normal, suggesting that ADCY5 is unlikely to link incretin signaling to cAMP generation and beta cell communication. Thus, ADCY5 preferentially affects glucose-

1 induced human islet dynamics, possibly through cAMP, which has been shown to increase GJ
2 conductance and trafficking [22,73,150], although this has only been so far demonstrated in
3 rodent tissues.

4
5 **ZnT8:** The R325W variant of SLC30A8, the gene encoding zinc transporter 8 (ZnT8), is
6 associated with reduced insulin secretion. ZnT8 is highly expressed in beta cells where its
7 activation leads to Zn^{2+} accumulation in secretory granules, promoting normal insulin
8 crystallization, storage and processing [151-154]. While global *ZnT8* deletion results in mild
9 insulin secretory deficits, which are only observed *in vivo* and are undetectable at the dispersed
10 islet level [151,152], beta-cell specific deletion of the same gene has been reported either to
11 inhibit [153] or stimulate [155] insulin release from isolated islets. Indeed, it has been suggested
12 that defects in glycemia resulting from either global or beta cell specific ZnT8 elimination
13 [152,153,155] are due to enhanced insulin clearance by the liver [155]. In any case, and
14 complicating the picture further, rare loss-of-function mutations in *SLC30A8* protect against
15 T2DM in man [156]. Nonetheless, alterations in ZnT8 expression lead to altered $\text{Ca}^{2+}/\text{Zn}^{2+}$ -
16 handling [133,152,157], and GJ gating is dependent on fine-regulation of both ions in the vicinity
17 of the plasma membrane [158,159]; whether this also applies to islets is unknown. Thus, while an
18 effect of *ZnT8* risk alleles on beta cell-beta cell connectivity is not entirely implausible, further
19 studies are required to assess effects of the gene on coordinated activity and the mechanisms
20 underlying this (*e.g.* changes in Cx36 expression or GJ function).

21
22 It should be noted that the studies concerning *ADCY5*, *TCF7L2* and beta cell connectivity were
23 conducted on models in which expression has essentially been eliminated (through gene silencing
24 or genomic deletion). It is likely that any phenotype observed *in vivo* in man is a consequence of
25 more subtle cellular changes coupled with exposure to a permissive environment. It remains to be
26 seen whether similar effects can be recapitulated in tissue obtained from normoglycemic donors
27 harboring specific risk alleles. Lastly, even the strongest GWAS hits only marginally contribute
28 to T2DM risk and effects of gene variants on beta cell coordination should not be overinterpreted
29 in the absence of defined mechanisms/targets.

30 31 **Rescuing beta cell connectivity during T2DM**

32 Since the intraislet regulation of insulin release may be altered by both genes and the environment
33 to reduce insulin secretion, beta cell connectivity may represent a novel target for the
34 pharmaceutical restoration of functional beta cell mass. While up-regulated GJ-signaling provides
35 a logical starting point for the enhancement of beta cell connectivity, investigation of Cx36-

modulating compounds has so far been complicated by their off-target effects. Notwithstanding, a recent study has described a panel of seventeen molecules that increase beta cell-beta cell communication, and further screening is warranted to validate their activity profiles and specificity [160]. In addition, atlases of both GPCR and paracrine factor expression/secretion have been reported for human and rodent islets [94,161], potentially accelerating the elucidation and development of putative candidates for manipulation of beta cell connectivity. Alternatively, personalized medicine/deep-phenotyping approaches [162] could be used to identify individuals where the beneficial effects of GLP-1 and GIP to enhance beta cell connectivity may be exploited [27,28]. For example, carriers of ADCY5 risk alleles are predicted to respond well to the insulin-raising actions of the incretins, as this gene preferentially impacts glucose action [148]. By contrast, obese subjects would potentially benefit more from the pro-communicatory effects of the sulfonylureas due to altered GLP-1 and GIP signaling inputs [27,163,164]

Future perspectives

The network description of beta cells is still in its infancy and more refined methods are required to better delineate connection topology. Without statistical methods such as Granger causality it is impossible to say whether coordinated behavior in an individual cell is the origin or consequence of the connections it shares with its neighbours [42,165]. Likewise, our understanding of the structural basis for functional connectivity is presently lacking and imaging approaches are required that allow the large-scale interrogation of any underlying physical cell-cell linkages. This is particularly applicable to human islets, where differences in architecture may lead to divergent regulation of insulin secretion and susceptibility to T2DM insults [28,37,64]. Lastly, it remains unknown how beta cell population dynamics are influenced by episodes of functional/pathological plasticity in the pancreas, and whether a wiring footprint persists during T2DM that can be exploited to restore insulin secretion.

Summary

The three-dimensional organization of beta cells into islets produces a gain-of-function in insulin release by fine-tuning beta cell intercommunication. Each islet operates as a self-supported signaling unit in which the spatiotemporally-precise propagation of information between neighboring and distant cell ensembles is facilitated by GJ, neural and paracrine communications. Using imaging approaches together with statistical methods borne from graph theory, the flow of information throughout the beta cell population can be monitored online and mapped. Pertinently, coordinated activity in rodent islets appears to be driven and orchestrated by a subpopulation of beta cells, and wiring density can be increased by both glucose and incretin to stimulate hormone

1 release. We therefore propose that, alongside “cell autonomous” effects, environmental and
2 genetic insults may target the intraislet regulation of insulin secretion to precipitate beta cell
3 dysfunction and glucose intolerance, contributing to the risk of developing T2DM.

4

1 REFERENCES

- 2 1. International Diabetes Federation. IDF Diabetes Atlas, 5th edn
- 3 2. Currie CJ, Poole CD, Gale EA (2009) The influence of glucose-lowering therapies on cancer risk in
- 4 type 2 diabetes. *Diabetologia* 52 (9):1766-1777. doi:10.1007/s00125-009-1440-6
- 5 3. Stitt AW (2010) AGEs and diabetic retinopathy. *Invest Ophthalmol Vis Sci* 51 (10):4867-4874.
- 6 doi:10.1167/iovs.10-5881
- 7 4. van de Bunt M, Gloyn AL (2012) A tale of two glucose transporters: how GLUT2 re-emerged as a
- 8 contender for glucose transport into the human beta cell. *Diabetologia* 55 (9):2312-2315.
- 9 doi:10.1007/s00125-012-2612-3
- 10 5. Iynedjian PB (1993) Mammalian glucokinase and its gene. *Biochem J* 293 (Pt 1):1-13
- 11 6. Prentki M, Matschinsky FM, Madiraju SR (2013) Metabolic signaling in fuel-induced insulin
- 12 secretion. *Cell Metab* 18 (2):162-185. doi:10.1016/j.cmet.2013.05.018
- 13 7. Rutter GA (2001) Nutrient-secretion coupling in the pancreatic islet beta-cell: recent advances. *Mol*
- 14 *Aspects Med* 22 (6):247-284
- 15 8. Tarasov AI, Semplici F, Ravier MA, Bellomo EA, Pullen TJ, Gilon P, Sekler I, Rizzuto R, Rutter
- 16 GA (2012) The mitochondrial Ca²⁺ uniporter MCU is essential for glucose-induced ATP increases in
- 17 pancreatic beta-cells. *PLoS One* 7 (7):e39722. doi:10.1371/journal.pone.0039722
- 18 9. Kennedy HJ, Pouli AE, Ainscow EK, Jouaville LS, Rizzuto R, Rutter GA (1999) Glucose generates
- 19 sub-plasma membrane ATP microdomains in single islet beta-cells. Potential role for strategically
- 20 located mitochondria. *The Journal of biological chemistry* 274 (19):13281-13291
- 21 10. Ashcroft FM, Harrison DE, Ashcroft SJ (1984) Glucose induces closure of single potassium
- 22 channels in isolated rat pancreatic beta-cells. *Nature* 312 (5993):446-448
- 23 11. Ashcroft FM, Gribble FM (1999) ATP-sensitive K⁺ channels and insulin secretion: their role in
- 24 health and disease. *Diabetologia* 42 (8):903-919. doi:10.1007/s001250051247
- 25 12. Ammala C, Ashcroft FM, Rorsman P (1993) Calcium-independent potentiation of insulin release
- 26 by cyclic AMP in single beta-cells. *Nature* 363 (6427):356-358. doi:10.1038/363356a0
- 27 13. Henquin JC (2000) Triggering and amplifying pathways of regulation of insulin secretion by
- 28 glucose. *Diabetes* 49 (11):1751-1760
- 29 14. Rutter GA, Tsuboi T, Ravier MA (2006) Ca²⁺ microdomains and the control of insulin secretion.
- 30 *Cell Calcium* 40 (5-6):539-551. doi:10.1016/j.ceca.2006.08.015
- 31 15. Emmanouilidou E, Teschemacher AG, Pouli AE, Nicholls LI, Seward EP, Rutter GA (1999)
- 32 Imaging Ca²⁺ concentration changes at the secretory vesicle surface with a recombinant targeted
- 33 cameleon. *Current biology : CB* 9 (16):915-918
- 34 16. Tsuboi T, Rutter GA (2003) Multiple forms of "kiss-and-run" exocytosis revealed by evanescent
- 35 wave microscopy. *Curr Biol* 13 (7):563-567
- 36 17. Rutter GA, Varadi A, Tsuboi T, Parton L, Ravier M (2006) Insulin secretion in health and disease:
- 37 genomics, proteomics and single vesicle dynamics. *Biochem Soc Trans* 34 (Pt 2):247-250.
- 38 doi:10.1042/BST20060247
- 39 18. Serre-Beinier V, Mas C, Calabrese A, Caton D, Bauquis J, Caille D, Charollais A, Cirulli V, Meda
- 40 P (2002) Connexins and secretion. *Biol Cell* 94 (7-8):477-492. doi:S0248490002000242 [pii]
- 41 19. Bavamian S, Klee P, Britan A, Populaire C, Caille D, Cancela J, Charollais A, Meda P (2007)
- 42 Islet-cell-to-cell communication as basis for normal insulin secretion. *Diabetes Obes Metab* 9 Suppl
- 43 2:118-132. doi:10.1111/j.1463-1326.2007.00780.x
- 44 20. Bosco D, Haefliger JA, Meda P (2011) Connexins: key mediators of endocrine function. *Physiol*
- 45 *Rev* 91 (4):1393-1445. doi:10.1152/physrev.00027.2010
- 46 21. Meda P (2013) Protein-mediated interactions of pancreatic islet cells. *Scientifica (Cairo)*
- 47 2013:621249. doi:10.1155/2013/621249
- 48 22. Farnsworth NL, Benninger RK (2014) New insights into the role of connexins in pancreatic islet
- 49 function and diabetes. *FEBS Lett* 588 (8):1278-1287. doi:10.1016/j.febslet.2014.02.035
- 50 23. Salomon D, Meda P (1986) Heterogeneity and contact-dependent regulation of hormone secretion
- 51 by individual B cells. *Exp Cell Res* 162 (2):507-520
- 52 24. Caton D, Calabrese A, Mas C, Serre-Beinier V, Wonkam A, Meda P (2002) Beta-cell crosstalk: a
- 53 further dimension in the stimulus-secretion coupling of glucose-induced insulin release. *Diabetes*
- 54 *Metab* 28 (6 Pt 2):3S45-53; discussion 43S108-112

25. Steiner DJ, Kim A, Miller K, Hara M (2010) Pancreatic islet plasticity: interspecies comparison of islet architecture and composition. *Islets* 2 (3):135-145
26. Hodson DJ, Mitchell RK, Marselli L, Pullen TJ, Brias SG, Semplici F, Everett KL, Cooper DM, Bugliani M, Marchetti P, Lavallard V, Bosco D, Piemonti L, Johnson PR, Hughes SJ, Li D, Li WH, Shapiro AM, Rutter GA (2014) ADCY5 couples glucose to insulin secretion in human islets. *Diabetes*. doi:10.2337/db13-1607
27. Hodson DJ, Mitchell RK, Bellomo EA, Sun G, Vinet L, Meda P, Li D, Li WH, Bugliani M, Marchetti P, Bosco D, Piemonti L, Johnson P, Hughes SJ, Rutter GA (2013) Lipotoxicity disrupts incretin-regulated human beta cell connectivity. *J Clin Invest* 123 (10):4182-4194. doi:10.1172/JCI68459
28. Rutter GA, Hodson DJ (2013) Minireview: intraislet regulation of insulin secretion in humans. *Mol Endocrinol* 27 (12):1984-1995. doi:10.1210/me.2013-1278
29. Stozar A, Gosak M, Dolensek J, Perc M, Marhl M, Rupnik MS, Korosak D (2013) Functional connectivity in islets of Langerhans from mouse pancreas tissue slices. *PLoS Comput Biol* 9 (2):e1002923. doi:10.1371/journal.pcbi.1002923
30. Stozar A, Dolensek J, Rupnik MS (2013) Glucose-stimulated calcium dynamics in islets of Langerhans in acute mouse pancreas tissue slices. *PLoS One* 8 (1):e54638. doi:10.1371/journal.pone.0054638
31. Benninger RK, Piston DW (2014) Cellular communication and heterogeneity in pancreatic islet insulin secretion dynamics. *Trends Endocrinol Metab*. doi:10.1016/j.tem.2014.02.005
32. Orci L, Unger RH (1975) Functional subdivision of islets of Langerhans and possible role of D cells. *Lancet* 2 (7947):1243-1244
33. Cabrera O, Berman DM, Kenyon NS, Ricordi C, Berggren PO, Caicedo A (2006) The unique cytoarchitecture of human pancreatic islets has implications for islet cell function. *Proc Natl Acad Sci U S A* 103 (7):2334-2339. doi:0510790103 [pii] 10.1073/pnas.0510790103
34. Nyman LR, Wells KS, Head WS, McCaughey M, Ford E, Brissova M, Piston DW, Powers AC (2008) Real-time, multidimensional in vivo imaging used to investigate blood flow in mouse pancreatic islets. *J Clin Invest* 118 (11):3790-3797. doi:10.1172/JCI36209
35. Cleaver O, Dor Y (2012) Vascular instruction of pancreas development. *Development* 139 (16):2833-2843. doi:10.1242/dev.065953
36. Brissova M, Fowler MJ, Nicholson WE, Chu A, Hirshberg B, Harlan DM, Powers AC (2005) Assessment of human pancreatic islet architecture and composition by laser scanning confocal microscopy. *J Histochem Cytochem* 53 (9):1087-1097. doi:jhc.5C6684.2005 [pii] 10.1369/jhc.5C6684.2005
37. Bosco D, Armanet M, Morel P, Niclauss N, Sgroi A, Muller YD, Giovannoni L, Parnaud G, Berney T (2010) Unique arrangement of alpha- and beta-cells in human islets of Langerhans. *Diabetes* 59 (5):1202-1210. doi:10.2337/db09-1177
38. Kilimnik G, Zhao B, Jo J, Periwai V, Witkowski P, Misawa R, Hara M (2011) Altered islet composition and disproportionate loss of large islets in patients with type 2 diabetes. *PLoS One* 6 (11):e27445. doi:10.1371/journal.pone.0027445
39. Halban PA, Wollheim CB, Blondel B, Meda P, Niesor EN, Mintz DH (1982) The possible importance of contact between pancreatic islet cells for the control of insulin release. *Endocrinology* 111 (1):86-94. doi:10.1210/endo-111-1-86
40. Hauge-Evans AC, Squires PE, Persaud SJ, Jones PM (1999) Pancreatic beta-cell-to-beta-cell interactions are required for integrated responses to nutrient stimuli: enhanced Ca²⁺ and insulin secretory responses of MIN6 pseudoislets. *Diabetes* 48 (7):1402-1408
41. Squires PE, Hauge-Evans AC, Persaud SJ, Jones PM (2000) Synchronization of Ca(2+)-signals within insulin-secreting pseudoislets: effects of gap-junctional uncouplers. *Cell Calcium* 27 (5):287-296. doi:10.1054/ceca.2000.0117
42. Hodson DJ, Molino F, Fontanaud P, Bonnefont X, Mollard P (2010) Investigating and Modelling Pituitary Endocrine Network Function. *J Neuroendocrinol* (22):1217-1225. doi:JNE2052 [pii] 10.1111/j.1365-2826.2010.02052.x

43. Takahashi N, Kishimoto T, Nemoto T, Kadowaki T, Kasai H (2002) Fusion pore dynamics and insulin granule exocytosis in the pancreatic islet. *Science* 297 (5585):1349-1352. doi:10.1126/science.1073806
44. Li D, Chen S, Bellomo EA, Tarasov AI, Kaut C, Rutter GA, Li WH (2011) Imaging dynamic insulin release using a fluorescent zinc indicator for monitoring induced exocytotic release (ZIMIR). *Proc Natl Acad Sci U S A* 108 (52):21063-21068. doi:10.1073/pnas.1109773109
45. Low JT, Mitchell JM, Do OH, Bax J, Rawlings A, Zavortink M, Morgan G, Parton RG, Gaisano HY, Thorn P (2013) Glucose principally regulates insulin secretion in mouse islets by controlling the numbers of granule fusion events per cell. *Diabetologia* 56 (12):2629-2637. doi:10.1007/s00125-013-3019-5
46. Pancholi J, Hodson DJ, Jobe K, Rutter GA, Goldup SM, Watkinson M (2014) Biologically targeted probes for Zn²⁺: a diversity oriented modular “click-SNAr-click” approach. *Chemical Science*. doi:10.1039/c4sc01249f
47. Akemann W, Mutoh H, Perron A, Rossier J, Knopfel T (2010) Imaging brain electric signals with genetically targeted voltage-sensitive fluorescent proteins. *Nat Methods* 7 (8):643-649. doi:10.1038/nmeth.1479
48. Hodson DJ, Romano N, Schaeffer M, Fontanaud P, Lafont C, Fiordelisio T, Mollard P (2012) Coordination of calcium signals by pituitary endocrine cells in situ. *Cell Calcium* 51 (3-4):222-230. doi:10.1016/j.ceca.2011.11.007
49. Dolensek J, Stozar A, Skelin Klemen M, Miller EW, Slak Rupnik M (2013) The relationship between membrane potential and calcium dynamics in glucose-stimulated beta cell syncytium in acute mouse pancreas tissue slices. *PLoS One* 8 (12):e82374. doi:10.1371/journal.pone.0082374
50. Schlegel W, Winiger BP, Mollard P, Vacher P, Wuarin F, Zahnd GR, Wollheim CB, Dufy B (1987) Oscillations of cytosolic Ca²⁺ in pituitary cells due to action potentials. *Nature* 329 (6141):719-721. doi:10.1038/329719a0
51. Peterlin ZA, Kozloski J, Mao BQ, Tsiola A, Yuste R (2000) Optical probing of neuronal circuits with calcium indicators. *Proc Natl Acad Sci U S A* 97 (7):3619-3624. doi:97/7/3619 [pii]
52. Cossart R, Ikegaya Y, Yuste R (2005) Calcium imaging of cortical networks dynamics. *Cell Calcium* 37 (5):451-457. doi:S0143-4160(05)00030-8 [pii] 10.1016/j.ceca.2005.01.013
53. Ikegaya Y, Le Bon-Jego M, Yuste R (2005) Large-scale imaging of cortical network activity with calcium indicators. *Neurosci Res* 52 (2):132-138. doi:S0168-0102(05)00063-5 [pii] 10.1016/j.neures.2005.02.004
54. Hodson DJ, Schaeffer M, Romano N, Fontanaud P, Lafont C, Birkenstock J, Molino F, Christian H, Lockey J, Carmignac D, Fernandez-Fuente M, Le Tissier P, Mollard P (2012) Existence of long-lasting experience-dependent plasticity in endocrine cell networks. *Nature Communications* 3:605. doi:10.1038/ncomms1612
55. Hodson DJ, Tarasov AI, Gimeno Brias S, Mitchell RK, Johnston NR, Haghollahi S, Cane MC, Bugliani M, Marchetti P, Bosco D, Johnson PR, Hughes SJ, Rutter GA (2014) Incretin-modulated beta cell energetics in intact islets of Langerhans. *Mol Endocrinol*:me20141038. doi:10.1210/me.2014-1038
56. Alon U (2007) Network motifs: theory and experimental approaches. *Nature Reviews Genetics* 8 (6):450-461. doi:10.1038/nrg2102
57. Bullmore E, Sporns O (2009) Complex brain networks: graph theoretical analysis of structural and functional systems. *Nat Rev Neurosci* 10 (3):186-198. doi:10.1038/nrn2575
58. Price DJ (1965) Networks of Scientific Papers. *Science* 149:510-515
59. Barabasi AL, Albert R (1999) Emergence of scaling in random networks. *Science* 286 (5439):509-512. doi:7898 [pii]
60. Barabasi AL (2009) Scale-free networks: a decade and beyond. *Science* 325 (5939):412-413. doi:325/5939/412 [pii] 10.1126/science.1173299
61. Bonifazi P, Goldin M, Picardo MA, Jorquera I, Cattani A, Bianconi G, Represa A, Ben-Ari Y, Cossart R (2009) GABAergic hub neurons orchestrate synchrony in developing hippocampal networks. *Science* 326 (5958):1419-1424. doi:326/5958/1419 [pii] 10.1126/science.1175509

62. Mollard P, Hodson DJ, Lafont C, Rizzoti K, Drouin J (2012) A tridimensional view of pituitary development and function. *Trends Endocrinol Metab.* doi:10.1016/j.tem.2012.02.004
63. Le Tissier PR, Hodson DJ, Lafont C, Fontanaud P, Schaeffer M, Mollard P (2012) Anterior pituitary cell networks. *Front Neuroendocrinol* 33 (3):252-266. doi:10.1016/j.yfrne.2012.08.002
64. Caicedo A (2013) Paracrine and autocrine interactions in the human islet: more than meets the eye. *Semin Cell Dev Biol* 24 (1):11-21. doi:10.1016/j.semcdb.2012.09.007
65. Serre-Beinier V, Le Gurun S, Belluardo N, Trovato-Salinaro A, Charollais A, Haeffliger JA, Condorelli DF, Meda P (2000) Cx36 preferentially connects beta-cells within pancreatic islets. *Diabetes* 49 (5):727-734
66. Condorelli DF, Belluardo N, Trovato-Salinaro A, Mudo G (2000) Expression of Cx36 in mammalian neurons. *Brain Res Brain Res Rev* 32 (1):72-85. doi:S0165017399000685 [pii]
67. Charpentier E, Cancela J, Meda P (2007) Beta cells preferentially exchange cationic molecules via connexin 36 gap junction channels. *Diabetologia* 50 (11):2332-2341. doi:10.1007/s00125-007-0807-9
68. Ravier MA, Guldenagel M, Charollais A, Gjinovci A, Caille D, Sohl G, Wollheim CB, Willecke K, Henquin JC, Meda P (2005) Loss of connexin36 channels alters beta-cell coupling, islet synchronization of glucose-induced Ca²⁺ and insulin oscillations, and basal insulin release. *Diabetes* 54 (6):1798-1807. doi:54/6/1798 [pii]
69. Speier S, Gjinovci A, Charollais A, Meda P, Rupnik M (2007) Cx36-mediated coupling reduces beta-cell heterogeneity, confines the stimulating glucose concentration range, and affects insulin release kinetics. *Diabetes* 56 (4):1078-1086. doi:10.2337/db06-0232
70. Rocheleau JV, Remedi MS, Granada B, Head WS, Koster JC, Nichols CG, Piston DW (2006) Critical role of gap junction coupled KATP channel activity for regulated insulin secretion. *PLoS Biology* 4 (2):e26. doi:05-PLBI-RA-0819R2 [pii]
- 10.1371/journal.pbio.0040026
71. Head WS, Orseth ML, Nunemaker CS, Satin LS, Piston DW, Benninger RK (2012) Connexin-36 gap junctions regulate in vivo first- and second-phase insulin secretion dynamics and glucose tolerance in the conscious mouse. *Diabetes* 61 (7):1700-1707. doi:10.2337/db11-1312
72. Serre-Beinier V, Bosco D, Zulianello L, Charollais A, Caille D, Charpentier E, Gauthier BR, Diaferia GR, Giepmans BN, Lupi R, Marchetti P, Deng S, Buhler L, Berney T, Cirulli V, Meda P (2009) Cx36 makes channels coupling human pancreatic beta-cells, and correlates with insulin expression. *Hum Mol Genet* 18 (3):428-439. doi:10.1093/hmg/ddn370
73. Farnsworth NL, Hemmati A, Pozzoli M, Benninger RK (2014) Fluorescence recovery after photobleaching reveals regulation and distribution of Cx36 gap junction coupling within mouse islets of langerhans. *J Physiol.* doi:10.1113/jphysiol.2014.276733
74. Ahren B (2000) Autonomic regulation of islet hormone secretion--implications for health and disease. *Diabetologia* 43 (4):393-410. doi:10.1007/s001250051322
75. Zhang M, Fendler B, Peercy B, Goel P, Bertram R, Sherman A, Satin L (2008) Long lasting synchronization of calcium oscillations by cholinergic stimulation in isolated pancreatic islets. *Biophys J* 95 (10):4676-4688. doi:S0006-3495(08)78607-7 [pii]
- 10.1529/biophysj.107.125088
76. Fendler B, Zhang M, Satin L, Bertram R (2009) Synchronization of pancreatic islet oscillations by intrapancreatic ganglia: a modeling study. *Biophys J* 97 (3):722-729. doi:10.1016/j.bpj.2009.05.016
77. Ahren B, Holst JJ (2001) The cephalic insulin response to meal ingestion in humans is dependent on both cholinergic and noncholinergic mechanisms and is important for postprandial glycemia. *Diabetes* 50 (5):1030-1038
78. Filipsson K, Kvist-Reimer M, Ahren B (2001) The neuropeptide pituitary adenylate cyclase-activating polypeptide and islet function. *Diabetes* 50 (9):1959-1969
79. Kurose T, Seino Y, Nishi S, Tsuji K, Taminato T, Tsuda K, Imura H (1990) Mechanism of sympathetic neural regulation of insulin, somatostatin, and glucagon secretion. *Am J Physiol* 258 (1 Pt 1):E220-227
80. Nilsson T, Arkhammar P, Rorsman P, Berggren PO (1988) Inhibition of glucose-stimulated insulin release by alpha 2-adrenoceptor activation is paralleled by both a repolarization and a reduction in cytoplasmic free Ca²⁺ concentration. *J Biol Chem* 263 (4):1855-1860

81. Kuo WN, Hodgins DS, Kuo JF (1973) Adenylate cyclase in islets of Langerhans. Isolation of islets and regulation of adenylate cyclase activity by various hormones and agents. *J Biol Chem* 248 (8):2705-2711
82. Asensio C, Jimenez M, Kuhne F, Rohner-Jeanrendaud F, Muzzin P (2005) The lack of beta-adrenoceptors results in enhanced insulin sensitivity in mice exhibiting increased adiposity and glucose intolerance. *Diabetes* 54 (12):3490-3495
83. Rodriguez-Diaz R, Abdulreda MH, Formoso AL, Gans I, Ricordi C, Berggren PO, Caicedo A (2011) Innervation patterns of autonomic axons in the human endocrine pancreas. *Cell Metabolism* 14 (1):45-54. doi:10.1016/j.cmet.2011.05.008
84. Rodriguez-Diaz R, Dando R, Jacques-Silva MC, Fachado A, Molina J, Abdulreda MH, Ricordi C, Roper SD, Berggren PO, Caicedo A (2011) Alpha cells secrete acetylcholine as a non-neuronal paracrine signal priming beta cell function in humans. *Nat Med* 17 (7):888-892. doi:10.1038/nm.2371
85. Molina J, Rodriguez-Diaz R, Fachado A, Jacques-Silva MC, Berggren PO, Caicedo A (2014) Control Of Insulin Secretion By Cholinergic Signaling In The Human Pancreatic Islet. *Diabetes*. doi:10.2337/db13-1371
86. Martin F, Soria B (1996) Glucose-induced $[Ca^{2+}]_i$ oscillations in single human pancreatic islets. *Cell Calcium* 20 (5):409-414
87. diIorio P, Rittenhouse AR, Bortell R, Jurczyk A (2014) Role of cilia in normal pancreas function and in diseased states. *Birth Defects Res C Embryo Today* 102 (2):126-138. doi:10.1002/bdrc.21064
88. Oh EC, Katsanis N (2012) Cilia in vertebrate development and disease. *Development* 139 (3):443-448. doi:10.1242/dev.050054
89. Cano DA, Sekine S, Hebrok M (2006) Primary cilia deletion in pancreatic epithelial cells results in cyst formation and pancreatitis. *Gastroenterology* 131 (6):1856-1869. doi:10.1053/j.gastro.2006.10.050
90. Ait-Lounis A, Baas D, Barras E, Benadiba C, Charollais A, Nlend Nlend R, Liegeois D, Meda P, Durand B, Reith W (2007) Novel function of the ciliogenic transcription factor RFX3 in development of the endocrine pancreas. *Diabetes* 56 (4):950-959. doi:10.2337/db06-1187
91. Granot Z, Swisa A, Magenheimer J, Stolovich-Rain M, Fujimoto W, Manduchi E, Miki T, Lennerz JK, Stoeckert CJ, Jr., Meyuhos O, Seino S, Permutt MA, Piwnicka-Worms H, Bardeesy N, Dor Y (2009) LKB1 regulates pancreatic beta cell size, polarity, and function. *Cell Metab* 10 (4):296-308. doi:10.1016/j.cmet.2009.08.010
92. Green JA, Mykityn K (2014) Neuronal primary cilia: an underappreciated signaling and sensory organelle in the brain. *Neuropsychopharmacology* 39 (1):244-245. doi:10.1038/npp.2013.203
93. Wang J, Silva M, Haas LA, Morsci NS, Nguyen KC, Hall DH, Barr MM (2014) C. elegans ciliated sensory neurons release extracellular vesicles that function in animal communication. *Curr Biol* 24 (5):519-525. doi:10.1016/j.cub.2014.01.002
94. Yang YH, Szabat M, Bragagnini C, Kott K, Helgason CD, Hoffman BG, Johnson JD (2011) Paracrine signalling loops in adult human and mouse pancreatic islets: netrins modulate beta cell apoptosis signalling via dependence receptors. *Diabetologia* 54 (4):828-842. doi:10.1007/s00125-010-2012-5
95. Ren J, Sherman A, Bertram R, Goforth PB, Nunemaker CS, Waters CD, Satin LS (2013) Slow oscillations of KATP conductance in mouse pancreatic islets provide support for electrical bursting driven by metabolic oscillations. *Am J Physiol Endocrinol Metab* 305 (7):E805-817. doi:10.1152/ajpendo.00046.2013
96. Merrins MJ, Fendler B, Zhang M, Sherman A, Bertram R, Satin LS (2010) Metabolic oscillations in pancreatic islets depend on the intracellular Ca^{2+} level but not Ca^{2+} oscillations. *Biophys J* 99 (1):76-84. doi:10.1016/j.bpj.2010.04.012
97. Nunemaker CS, Satin LS (2004) Comparison of metabolic oscillations from mouse pancreatic beta cells and islets. *Endocrine* 25 (1):61-67. doi:ENDO:25:1:61 [pii] 10.1385/ENDO:25:1:61
98. Li J, Shuai HY, Gylfe E, Tengholm A (2013) Oscillations of sub-membrane ATP in glucose-stimulated beta cells depend on negative feedback from Ca^{2+} . *Diabetologia* 56 (7):1577-1586. doi:10.1007/s00125-013-2894-0
99. Bennett BD, Jetton TL, Ying G, Magnuson MA, Piston DW (1996) Quantitative subcellular imaging of glucose metabolism within intact pancreatic islets. *J Biol Chem* 271 (7):3647-3651

100. Piston DW, Knobel SM (1999) Quantitative imaging of metabolism by two-photon excitation microscopy. *Methods Enzymol* 307:351-368
101. Nunemaker CS, Dishinger JF, Dula SB, Wu R, Merrins MJ, Reid KR, Sherman A, Kennedy RT, Satin LS (2009) Glucose metabolism, islet architecture, and genetic homogeneity in imprinting of [Ca²⁺]_i and insulin rhythms in mouse islets. *PLoS One* 4 (12):e8428. doi:10.1371/journal.pone.0008428
102. Kohen E, Kohen C, Thorell B, Mintz DH, Rabinovitch A (1979) Intercellular communication in pancreatic islet monolayer cultures: a microfluorometric study. *Science* 204 (4395):862-865
103. Meda P, Amherdt M, Perrelet A, Orci L (1981) Metabolic coupling between cultured pancreatic b-cells. *Exp Cell Res* 133 (2):421-430
104. Reimann F, Gribble FM (2002) Glucose-sensing in glucagon-like peptide-1-secreting cells. *Diabetes* 51 (9):2757-2763
105. Tolhurst G, Reimann F, Gribble FM (2009) Nutritional regulation of glucagon-like peptide-1 secretion. *J Physiol* 587 (Pt 1):27-32. doi:10.1113/jphysiol.2008.164012
106. Parker HE, Wallis K, le Roux CW, Wong KY, Reimann F, Gribble FM (2012) Molecular mechanisms underlying bile acid-stimulated glucagon-like peptide-1 secretion. *Br J Pharmacol* 165 (2):414-423. doi:10.1111/j.1476-5381.2011.01561.x
107. Gomez E, Pritchard C, Herbert TP (2002) cAMP-dependent protein kinase and Ca²⁺ influx through L-type voltage-gated calcium channels mediate Raf-independent activation of extracellular regulated kinase in response to glucagon-like peptide-1 in pancreatic beta-cells. *J Biol Chem* 277 (50):48146-48151. doi:10.1074/jbc.M209165200
108. Leech CA, Dzura I, Chepurny OG, Kang G, Schwede F, Genieser HG, Holz GG (2011) Molecular physiology of glucagon-like peptide-1 insulin secretagogue action in pancreatic beta cells. *Prog Biophys Mol Biol* 107 (2):236-247. doi:10.1016/j.pbiomolbio.2011.07.005
109. Ravier MA, Leduc M, Richard J, Linck N, Varrault A, Pirot N, Roussel MM, Bockaert J, Dalle S, Bertrand G (2013) beta-Arrestin2 plays a key role in the modulation of the pancreatic beta cell mass in mice. *Diabetologia*. doi:10.1007/s00125-013-3130-7
110. Tsuboi T, da Silva Xavier G, Holz GG, Jouaville LS, Thomas AP, Rutter GA (2003) Glucagon-like peptide-1 mobilizes intracellular Ca²⁺ and stimulates mitochondrial ATP synthesis in pancreatic MIN6 beta-cells. *The Biochemical journal* 369 (Pt 2):287-299. doi:10.1042/BJ20021288
111. Peyot ML, Gray JP, Lamontagne J, Smith PJ, Holz GG, Madiraju SR, Prentki M, Heart E (2009) Glucagon-like peptide-1 induced signaling and insulin secretion do not drive fuel and energy metabolism in primary rodent pancreatic beta-cells. *PLoS One* 4 (7):e6221. doi:10.1371/journal.pone.0006221
112. Berg J, Hung YP, Yellen G (2009) A genetically encoded fluorescent reporter of ATP:ADP ratio. *Nat Methods* 6 (2):161-166. doi:10.1038/nmeth.1288
113. Hodson DJ, Mitchell RK, Johnston N, Thorens B, Ferrer J, Rutter GA (2014) Optical control of beta cell function. *Diabet Med* 31:6-6
114. Nauck MA, Homberger E, Siegel EG, Allen RC, Eaton RP, Ebert R, Creutzfeldt W (1986) Incretin effects of increasing glucose loads in man calculated from venous insulin and C-peptide responses. *J Clin Endocrinol Metab* 63 (2):492-498. doi:10.1210/jcem-63-2-492
115. Nauck MA, Heimesaat MM, Orskov C, Holst JJ, Ebert R, Creutzfeldt W (1993) Preserved incretin activity of glucagon-like peptide 1 [7-36 amide] but not of synthetic human gastric inhibitory polypeptide in patients with type-2 diabetes mellitus. *J Clin Invest* 91 (1):301-307. doi:10.1172/JCI116186
116. Kjems LL, Holst JJ, Volund A, Madsbad S (2003) The influence of GLP-1 on glucose-stimulated insulin secretion: effects on beta-cell sensitivity in type 2 and nondiabetic subjects. *Diabetes* 52 (2):380-386
117. Meier JJ, Nauck MA (2010) Is the diminished incretin effect in type 2 diabetes just an epiphenomenon of impaired beta-cell function? *Diabetes* 59 (5):1117-1125. doi:10.2337/db09-1899
118. Muscelli E, Mari A, Casolaro A, Camastra S, Seghieri G, Gastaldelli A, Holst JJ, Ferrannini E (2008) Separate impact of obesity and glucose tolerance on the incretin effect in normal subjects and type 2 diabetic patients. *Diabetes* 57 (5):1340-1348. doi:10.2337/db07-1315
119. Knop FK, Aaboe K, Vilsboll T, Volund A, Holst JJ, Krarup T, Madsbad S (2012) Impaired incretin effect and fasting hyperglucagonaemia characterizing type 2 diabetic subjects are early signs

of dysmetabolism in obesity. *Diabetes Obes Metab* 14 (6):500-510. doi:10.1111/j.1463-1326.2011.01549.x

120. Allagnat F, Alonso F, Martin D, Abderrahmani A, Waeber G, Haefliger JA (2008) ICER-1gamma overexpression drives palmitate-mediated connexin36 down-regulation in insulin-secreting cells. *J Biol Chem* 283 (9):5226-5234. doi:10.1074/jbc.M708181200

121. Haefliger JA, Martin D, Favre D, Petremand Y, Mazzolai L, Abderrahmani A, Meda P, Waeber G, Allagnat F (2013) Reduction of connexin36 content by ICER-1 contributes to insulin-secreting cells apoptosis induced by oxidized LDL particles. *PLoS One* 8 (1):e55198. doi:10.1371/journal.pone.0055198

122. Newman B, Selby JV, King MC, Slemenda C, Fabsitz R, Friedman GD (1987) Concordance for type 2 (non-insulin-dependent) diabetes mellitus in male twins. *Diabetologia* 30 (10):763-768

123. Pierce M, Keen H, Bradley C (1995) Risk of Diabetes in Offspring of Parents with Non-insulin-dependent Diabetes. *Diabet Med* 12 (1):6-13. doi:10.1111/j.1464-5491.1995.tb02054.x

124. Medici F, Hawa M, Ianari A, Pyke DA, Leslie RD (1999) Concordance rate for type II diabetes mellitus in monozygotic twins: actuarial analysis. *Diabetologia* 42 (2):146-150. doi:10.1007/s001250051132

125. Rutter GA (2014) Understanding genes identified by genome-wide association studies for Type 2 diabetes. *Diabet Med*. doi:10.1111/dme.12579

126. McCarthy MI (2010) Genomics, type 2 diabetes, and obesity. *N Engl J Med* 363 (24):2339-2350. doi:10.1056/NEJMra0906948

127. Lyssenko V, Lupi R, Marchetti P, Del Guerra S, Orho-Melander M, Almgren P, Sjogren M, Ling C, Eriksson KF, Lethagen AL, Mancarella R, Berglund G, Tuomi T, Nilsson P, Del Prato S, Groop L (2007) Mechanisms by which common variants in the TCF7L2 gene increase risk of type 2 diabetes. *J Clin Invest* 117 (8):2155-2163. doi:10.1172/JCI30706

128. Salonen JT, Uimari P, Aalto JM, Pirskanen M, Kaikkonen J, Todorova B, Hypponen J, Korhonen VP, Asikainen J, Devine C, Tuomainen TP, Luedemann J, Nauck M, Kerner W, Stephens RH, New JP, Ollier WE, Gibson JM, Payton A, Horan MA, Pendleton N, Mahoney W, Meyre D, Delplanque J, Froguel P, Luzzatto O, Yakir B, Darvasi A (2007) Type 2 diabetes whole-genome association study in four populations: the DiaGen consortium. *Am J Hum Genet* 81 (2):338-345. doi:10.1086/520599

129. Vaxillaire M, Veslot J, Dina C, Proenca C, Cauchi S, Charpentier G, Tichet J, Fumeron F, Marre M, Meyre D, Balkau B, Froguel P (2008) Impact of common type 2 diabetes risk polymorphisms in the DESIR prospective study. *Diabetes* 57 (1):244-254. doi:10.2337/db07-0615

130. Palmer ND, Lehtinen AB, Langefeld CD, Campbell JK, Haffner SM, Norris JM, Bergman RN, Goodarzi MO, Rotter JJ, Bowden DW (2008) Association of TCF7L2 gene polymorphisms with reduced acute insulin response in Hispanic Americans. *J Clin Endocrinol Metab* 93 (1):304-309. doi:10.1210/jc.2007-1225

131. Schafer SA, Tschritter O, Machicao F, Thamer C, Stefan N, Gallwitz B, Holst JJ, Dekker JM, Hart LM, Nijpels G, van Haeften TW, Haring HU, Fritsche A (2007) Impaired glucagon-like peptide-1-induced insulin secretion in carriers of transcription factor 7-like 2 (TCF7L2) gene polymorphisms. *Diabetologia* 50 (12):2443-2450. doi:10.1007/s00125-007-0753-6

132. Pilgaard K, Jensen CB, Schou JH, Lyssenko V, Wegner L, Brons C, Vilsboll T, Hansen T, Madsbad S, Holst JJ, Volund A, Poulsen P, Groop L, Pedersen O, Vaag AA (2009) The T allele of rs7903146 TCF7L2 is associated with impaired insulinotropic action of incretin hormones, reduced 24 h profiles of plasma insulin and glucagon, and increased hepatic glucose production in young healthy men. *Diabetologia* 52 (7):1298-1307. doi:10.1007/s00125-009-1307-x

133. da Silva Xavier G, Loder MK, McDonald A, Tarasov AI, Carzaniga R, Kronenberger K, Barg S, Rutter GA (2009) TCF7L2 regulates late events in insulin secretion from pancreatic islet beta-cells. *Diabetes* 58 (4):894-905. doi:10.2337/db08-1187

134. Shu L, Sauter NS, Schulthess FT, Matveyenko AV, Oberholzer J, Maedler K (2008) Transcription factor 7-like 2 regulates beta-cell survival and function in human pancreatic islets. *Diabetes* 57 (3):645-653. doi:10.2337/db07-0847

135. da Silva Xavier G, Mondragon A, Sun G, Chen L, McGinty JA, French PM, Rutter GA (2012) Abnormal glucose tolerance and insulin secretion in pancreas-specific Tcf7l2-null mice. *Diabetologia* 55 (10):2667-2676. doi:10.1007/s00125-012-2600-7

136. da Silva Xavier G, Mondragon A, Mitchell RK, Hodson DJ, Ferrer J, Thoren B, Chen L, McGinty JA, French PM, Rutter GA Defective glucose homeostasis in mice inactivated selectively for Tcf7l2 in the adult beta cell with an Ins1-controlled Cre. In: EASD, Vienna, 2014.
137. Boj SF, van Es JH, Huch M, Li VS, Jose A, Hatzis P, Mokry M, Haegebarth A, van den Born M, Chambon P, Voshol P, Dor Y, Cuppen E, Fillat C, Clevers H (2012) Diabetes Risk Gene and Wnt Effector Tcf7l2/TCF4 Controls Hepatic Response to Perinatal and Adult Metabolic Demand. *Cell* 151 (7):1595-1607. doi:10.1016/j.cell.2012.10.053
138. Shu L, Matveyenko AV, Kerr-Conte J, Cho JH, McIntosh CH, Maedler K (2009) Decreased TCF7L2 protein levels in type 2 diabetes mellitus correlate with downregulation of GIP- and GLP-1 receptors and impaired beta-cell function. *Hum Mol Genet* 18 (13):2388-2399. doi:10.1093/hmg/ddp178
139. Rosengren AH, Braun M, Mahdi T, Andersson SA, Travers ME, Shigeto M, Zhang E, Almgren P, Ladvall C, Axelsson AS, Edlund A, Pedersen MG, Jonsson A, Ramacheya R, Tang Y, Walker JN, Barrett A, Johnson PR, Lyssenko V, McCarthy MI, Groop L, Salehi A, Gloyn AL, Renstrom E, Rorsman P, Eliasson L (2012) Reduced insulin exocytosis in human pancreatic beta-cells with gene variants linked to type 2 diabetes. *Diabetes* 61 (7):1726-1733. doi:10.2337/db11-1516
140. Loder MK, da Silva Xavier G, McDonald A, Rutter GA (2008) TCF7L2 controls insulin gene expression and insulin secretion in mature pancreatic beta-cells. *Biochem Soc Trans* 36 (Pt 3):357-359. doi:10.1042/BST0360357
141. Zhou Y, Park SY, Su J, Bailey K, Ottosson-Laakso E, Shcherbina L, Oskolkov N, Zhang E, Thevenin T, Fadista J, Bennet H, Vikman P, Wierup N, Fex M, Rung J, Wollheim C, Nobrega M, Renstrom E, Groop L, Hansson O (2014) TCF7L2 is a master regulator of insulin production and processing. *Hum Mol Genet*. doi:10.1093/hmg/ddu359
142. Yoshimura M, Cooper DM (1992) Cloning and expression of a Ca(2+)-inhibitable adenylyl cyclase from NCB-20 cells. *Proc Natl Acad Sci U S A* 89 (15):6716-6720
143. Yan L, Vatner DE, O'Connor JP, Ivessa A, Ge H, Chen W, Hirotani S, Ishikawa Y, Sadoshima J, Vatner SF (2007) Type 5 adenylyl cyclase disruption increases longevity and protects against stress. *Cell* 130 (2):247-258. doi:10.1016/j.cell.2007.05.038
144. Holz GG, Leech CA, Chepurny OG (2014) New insights concerning the molecular basis for defective glucoregulation in soluble adenylyl cyclase knockout mice. *Biochim Biophys Acta*. doi:10.1016/j.bbadis.2014.06.023
145. Leech CA, Castonguay MA, Habener JF (1999) Expression of adenylyl cyclase subtypes in pancreatic beta-cells. *Biochem Biophys Res Commun* 254 (3):703-706. doi:10.1006/bbrc.1998.9906
146. Dupuis J, Langenberg C, Prokopenko I, Saxena R, Soranzo N, Jackson AU, Wheeler E, Glazer NL, Bouatia-Naji N, Gloyn AL, Lindgren CM, Magi R, Morris AP, Randall J, Johnson T, Elliott P, Rybin D, Thorleifsson G, Steinthorsdottir V, Henneman P, Grallert H, Dehghan A, Hottenga JJ, Franklin CS, Navarro P, Song K, Goel A, Perry JR, Egan JM, Lajunen T, Grarup N, Sparso T, Doney A, Voight BF, Stringham HM, Li M, Kanoni S, Shrader P, Cavalcanti-Proenca C, Kumari M, Qi L, Timpson NJ, Gieger C, Zabena C, Rocheleau G, Ingelsson E, An P, O'Connell J, Luan J, Elliott A, McCarroll SA, Payne F, Roccascaccia RM, Pattou F, Sethupathy P, Ardlie K, Ariyurek Y, Balkau B, Barter P, Beilby JP, Ben-Shlomo Y, Benediktsson R, Bennett AJ, Bergmann S, Bochud M, Boerwinkle E, Bonnefond A, Bonnycastle LL, Borch-Johnsen K, Bottcher Y, Brunner E, Bumpstead SJ, Charpentier G, Chen YD, Chines P, Clarke R, Coin LJ, Cooper MN, Cornelis M, Crawford G, Crisponi L, Day IN, de Geus EJ, Delplanque J, Dina C, Erdos MR, Fedson AC, Fischer-Rosinsky A, Forouhi NG, Fox CS, Frants R, Franzosi MG, Galan P, Goodarzi MO, Graessler J, Groves CJ, Grundy S, Gwilliam R, Gyllenstein U, Hadjadj S, Hallmans G, Hammond N, Han X, Hartikainen AL, Hassanali N, Hayward C, Heath SC, Hercberg S, Herder C, Hicks AA, Hillman DR, Hingorani AD, Hofman A, Hui J, Hung J, Isomaa B, Johnson PR, Jorgensen T, Jula A, Kaakinen M, Kaprio J, Kesaniemi YA, Kivimaki M, Knight B, Koskinen S, Kovacs P, Kyvik KO, Lathrop GM, Lawlor DA, Le Bacquer O, Lecoeur C, Li Y, Lyssenko V, Mahley R, Mangino M, Manning AK, Martinez-Larrad MT, McAteer JB, McCulloch LJ, McPherson R, Meisinger C, Melzer D, Meyre D, Mitchell BD, Morken MA, Mukherjee S, Naitza S, Narisu N, Neville MJ, Oostra BA, Orru M, Pakyz R, Palmer CN, Paolisso G, Pattaro C, Pearson D, Peden JF, Pedersen NL, Perola M, Pfeiffer AF, Pichler I, Polasek O, Posthuma D, Potter SC, Pouta A, Province MA, Psaty BM, Rathmann W, Rayner NW,

Rice K, Ripatti S, Rivadeneira F, Roden M, Rolandsson O, Sandbaek A, Sandhu M, Sanna S, Sayer AA, Scheet P, Scott LJ, Seedorf U, Sharp SJ, Shields B, Sigurethsson G, Sijbrands EJ, Silveira A, Simpson L, Singleton A, Smith NL, Sovio U, Swift A, Syddall H, Syvanen AC, Tanaka T, Thorand B, Tichet J, Tonjes A, Tuomi T, Uitterlinden AG, van Dijk KW, van Hoek M, Varma D, Visvikis-Siest S, Vitart V, Vogelzangs N, Waeber G, Wagner PJ, Walley A, Walters GB, Ward KL, Watkins H, Weedon MN, Wild SH, Willemsen G, Witteman JC, Yarnell JW, Zeggini E, Zelenika D, Zethelius B, Zhai G, Zhao JH, Zillikens MC, Borecki IB, Loos RJ, Meneton P, Magnusson PK, Nathan DM, Williams GH, Hattersley AT, Silander K, Salomaa V, Smith GD, Bornstein SR, Schwarz P, Spranger J, Karpe F, Shuldiner AR, Cooper C, Dedoussis GV, Serrano-Rios M, Morris AD, Lind L, Palmer LJ, Hu FB, Franks PW, Ebrahim S, Marmot M, Kao WH, Pankow JS, Sampson MJ, Kuusisto J, Laakso M, Hansen T, Pedersen O, Pramstaller PP, Wichmann HE, Illig T, Rudan I, Wright AF, Stumvoll M, Campbell H, Wilson JF, Bergman RN, Buchanan TA, Collins FS, Mohlke KL, Tuomilehto J, Valle TT, Altshuler D, Rotter JI, Siscovick DS, Penninx BW, Boomsma DI, Deloukas P, Spector TD, Frayling TM, Ferrucci L, Kong A, Thorsteinsdottir U, Stefansson K, van Duijn CM, Aulchenko YS, Cao A, Scuteri A, Schlessinger D, Uda M, Ruokonen A, Jarvelin MR, Waterworth DM, Vollenweider P, Peltonen L, Mooser V, Abecasis GR, Wareham NJ, Sladek R, Froguel P, Watanabe RM, Meigs JB, Groop L, Boehnke M, McCarthy MI, Florez JC, Barroso I (2010) New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nat Genet* 42 (2):105-116. doi:ng.520 [pii] 10.1038/ng.520

147. Rees SD, Hydrie MZ, O'Hare JP, Kumar S, Shera AS, Basit A, Barnett AH, Kelly MA (2011) Effects of 16 genetic variants on fasting glucose and type 2 diabetes in South Asians: ADCY5 and GLIS3 variants may predispose to type 2 diabetes. *PLoS One* 6 (9):e24710. doi:10.1371/journal.pone.0024710

148. Hodson DJ, Mitchell RK, Marselli L, Pullen TJ, Gimeno Brias S, Semplici F, Everett KL, Cooper DMF, Bugliani M, Marchetti P, Lavallard V, Bosco D, Piemonti L, Johnson PR, Hughes SJ, Li D, Li W-H, Shapiro AMJ, Rutter GA (2014) ADCY5 couples glucose to insulin secretion in human islets *Diabetes* In press

149. Wagner R, Dudziak K, Herzberg-Schafer SA, Machicao F, Stefan N, Staiger H, Haring HU, Fritsche A (2011) Glucose-raising genetic variants in MADD and ADCY5 impair conversion of proinsulin to insulin. *PLoS One* 6 (8):e23639. doi:10.1371/journal.pone.0023639

150. Mears D, Sheppard NF, Jr., Atwater I, Rojas E (1995) Magnitude and modulation of pancreatic beta-cell gap junction electrical conductance in situ. *J Membr Biol* 146 (2):163-176

151. Lemaire K, Ravier MA, Schraenen A, Creemers JW, Van de Plas R, Granvik M, Van Lommel L, Waelkens E, Chimienti F, Rutter GA, Gilon P, in't Veld PA, Schuit FC (2009) Insulin crystallization depends on zinc transporter ZnT8 expression, but is not required for normal glucose homeostasis in mice. *Proc Natl Acad Sci U S A* 106 (35):14872-14877. doi:10.1073/pnas.0906587106

152. Nicolson TJ, Bellomo EA, Wijesekara N, Loder MK, Baldwin JM, Gyulkhandanyan AV, Koshkin V, Tarasov AI, Carzaniga R, Kronenberger K, Taneja TK, da Silva Xavier G, Libert S, Froguel P, Scharfmann R, Stetsyuk V, Ravassard P, Parker H, Gribble FM, Reimann F, Sladek R, Hughes SJ, Johnson PR, Masseboeuf M, Burcelin R, Baldwin SA, Liu M, Lara-Lemus R, Arvan P, Schuit FC, Wheeler MB, Chimienti F, Rutter GA (2009) Insulin storage and glucose homeostasis in mice null for the granule zinc transporter ZnT8 and studies of the type 2 diabetes-associated variants. *Diabetes* 58 (9):2070-2083. doi:10.2337/db09-0551

153. Wijesekara N, Dai FF, Hardy AB, Giglou PR, Bhattacharjee A, Koshkin V, Chimienti F, Gaisano HY, Rutter GA, Wheeler MB (2010) Beta cell-specific Znt8 deletion in mice causes marked defects in insulin processing, crystallisation and secretion. *Diabetologia* 53 (8):1656-1668. doi:10.1007/s00125-010-1733-9

154. Rutter GA (2010) Think zinc: New roles for zinc in the control of insulin secretion. *Islets* 2 (1):49-50. doi:10.4161/isl.2.1.10259

155. Tamaki M, Fujitani Y, Hara A, Uchida T, Tamura Y, Takeno K, Kawaguchi M, Watanabe T, Ogiwara T, Fukunaka A, Shimizu T, Mita T, Kanazawa A, Imaizumi MO, Abe T, Kiyonari H, Hojyo S, Fukada T, Kawachi T, Nagamatsu S, Hirano T, Kawamori R, Watada H (2013) The diabetes-susceptible gene SLC30A8/ZnT8 regulates hepatic insulin clearance. *J Clin Invest* 123 (10):4513-4524. doi:10.1172/JCI68807

156. Flannick J, Thorleifsson G, Beer NL, Jacobs SB, Grarup N, Burt NP, Mahajan A, Fuchsberger C, Atzmon G, Benediktsson R, Blangero J, Bowden DW, Brandslund I, Brosnan J, Burslem F, Chambers J, Cho YS, Christensen C, Douglas DA, Duggirala R, Dymek Z, Farjoun Y, Fennell T, Fontanillas P, Forsen T, Gabriel S, Glaser B, Gudbjartsson DF, Hanis C, Hansen T, Hreidarsson AB, Hveem K, Ingelsson E, Isomaa B, Johansson S, Jorgensen T, Jorgensen ME, Kathiresan S, Kong A, Kooner J, Kravic J, Laakso M, Lee JY, Lind L, Lindgren CM, Linneberg A, Masson G, Meitinger T, Mohlke KL, Molven A, Morris AP, Potluri S, Rauramaa R, Ribel-Madsen R, Richard AM, Rolph T, Salomaa V, Segre AV, Skarstrand H, Steinthorsdottir V, Stringham HM, Sulem P, Tai ES, Teo YY, Teslovich T, Thorsteinsdottir U, Trimmer JK, Tuomi T, Tuomilehto J, Vaziri-Sani F, Voight BF, Wilson JG, Boehnke M, McCarthy MI, Njolstad PR, Pedersen O, Groop L, Cox DR, Stefansson K, Altshuler D (2014) Loss-of-function mutations in SLC30A8 protect against type 2 diabetes. *Nat Genet* 46 (4):357-363. doi:10.1038/ng.2915
157. Gerber PA, Bellomo EA, Hodson DJ, Meur G, Solomou A, Mitchell RK, Hollinshead M, Chimienti F, Bosco D, Hughes SJ, Johnson PR, Rutter GA (2014) Hypoxia lowers SLC30A8/ZnT8 expression and free cytosolic Zn²⁺ in pancreatic beta cells. *Diabetologia*. doi:10.1007/s00125-014-3266-0
158. Sun Z, Zhang DQ, McMahon DG (2009) Zinc modulation of hemi-gap-junction channel currents in retinal horizontal cells. *J Neurophysiol* 101 (4):1774-1780. doi:90581.2008 [pii] 10.1152/jn.90581.2008
159. Lurtz MM, Louis CF (2007) Intracellular calcium regulation of connexin43. *Am J Physiol Cell Physiol* 293 (6):C1806-1813. doi:00630.2006 [pii] 10.1152/ajpcell.00630.2006
160. Bavamian S, Pontes H, Cancela J, Charollais A, Startchik S, Van de Ville D, Meda P (2012) The intercellular synchronization of Ca²⁺ oscillations evaluates Cx36-dependent coupling. *PLoS One* 7 (7):e41535. doi:10.1371/journal.pone.0041535
161. Amisten S, Salehi A, Rorsman P, Jones PM, Persaud SJ (2013) An atlas and functional analysis of G-protein coupled receptors in human islets of Langerhans. *Pharmacol Ther* 139 (3):359-391. doi:10.1016/j.pharmthera.2013.05.004
162. Zhou K, Pearson ER (2013) Insights from genome-wide association studies of drug response. *Annu Rev Pharmacol Toxicol* 53:299-310. doi:10.1146/annurev-pharmtox-011112-140237
163. Meda P, Michaels RL, Halban PA, Orci L, Sheridan JD (1983) In vivo modulation of gap junctions and dye coupling between B-cells of the intact pancreatic islet. *Diabetes* 32 (9):858-868
164. Donnelly LA, Doney ASF, Hattersley AT, Morris AD, Pearson ER (2006) The effect of obesity on glycaemic response to metformin or sulphonylureas in Type 2 diabetes. *Diabet Med* 23 (2):128-133. doi:DOI 10.1111/j.1464-5491.2005.01755.x
165. Kim S, Putrino D, Ghosh S, Brown EN (2011) A Granger causality measure for point process models of ensemble neural spiking activity. *PLoS Comput Biol* 7 (3):e1001110. doi:10.1371/journal.pcbi.1001110

Acknowledgements

The writing of this review article was supported by a Diabetes UK R.D. Lawrence Research Fellowship (12/0004431) to D.J.H. and Wellcome Trust Senior Investigator (WT098424AIA), MRC Programme (MR/J0003042/1), Diabetes UK Project Grant (11/0004210) and Royal Society Wolfson Research Merit Awards to G.A.R. The work leading to this publication has received support from the Innovative Medicines Initiative Joint Undertaking under grant agreement n° 155005 (IMIDIA), resources of which are composed of financial contribution from the European Union's Seventh Framework Programme (FP7/2007-2013) and EFPIA companies' in kind contribution (G.A.R.). Lastly, we thank Miss Maria Paiva Pessoa for assistance with the figures.

FIGURE LEGENDS

Figure 1: Imaging and mapping beta cell network topology. (Above) Functional multicellular Ca^{2+} imaging is used to monitor the large-scale organization of glucose-induced population dynamics (*above, left*). By subjecting the resulting traces (from ~ 50-100 individual cells per islet) to correlation analyses, cells with coordinated activity can be identified and a functional connectivity map plotted based upon position within the imaged field (x-y) (*above, right*). Scale-free connection distributions are typified by a minority of cells that host the majority of connections (nodes), while maintaining streamlined information flow due to a short pathlength. Although robust in the face of random attack, they are prone to collapse following a targeted attack (*below, left*). By contrast, non-scale free networks (*e.g.* random or lattice) may not efficiently propagate signals due to a long pathlength, and random attacks significantly reduce capacity (*below, right*).

Figure 2: Schematic showing single cell and population level beta cell signaling. At the molecular level, glucose is transported into the beta cell before undergoing glycolysis to increase the ratio of free cytosolic ATP:ADP. This closes K_{ATP} channels, leading to opening of VDCC, Ca^{2+} influx and Ca^{2+} -dependent exocytosis. At the population level, beta cell dynamics are further dictated by signaling circuits involving paracrine, juxtacrine, autocrine, electrotonic (GJ), neural and ciliary communications.

Figure 3: Potential mechanisms by which T2D-associated genes may alter beta cell connectivity. *ZnT8* gene variants disrupt cytosolic Ca^{2+} and Zn^{2+} handling, and both of these ions are required for normal GJ activity. *ADCY5* gene variants decrease glucose-stimulated cAMP rises, a second messenger shown to increase GJ communications between beta cells. By contrast, *TCF7L2* gene variants may disrupt normal GJ function through effects upon glucose-stimulated Ca^{2+} increases, as well as GLP-1-stimulated cAMP generation.